

Genetic structure is stronger across human-impacted habitats than among islands in the coral *Porites lobata*

Kaho H. Tisthammer^{1,2}, Zac H. Forsman³, Robert J. Toonen³ and Robert H. Richmond¹

¹Kewalo Marine Laboratory, University of Hawaii at Manoa, Honolulu, HI, United States of America

²Department of Biology, San Francisco State University, San Francisco, CA, United States of America

³Hawaii Institute of Marine Biology, University of Hawaii at Manoa, Kaneohe, HI, United States of America

ABSTRACT

We examined genetic structure in the lobe coral *Porites lobata* among pairs of highly variable and high-stress nearshore sites and adjacent less variable and less impacted offshore sites on the islands of O'ahu and Maui, Hawai'i. Using an analysis of molecular variance framework, we tested whether populations were more structured by geographic distance or environmental extremes. The genetic patterns we observed followed isolation by environment, where nearshore and adjacent offshore populations showed significant genetic structure at both locations (AMOVA $F_{ST} = 0.04\sim 0.19$, $P < 0.001$), but no significant isolation by distance between islands. Strikingly, corals from the two nearshore sites with higher levels of environmental stressors on different islands over 100 km apart with similar environmentally stressful conditions were genetically closer ($F_{ST} = 0.0$, $P = 0.73$) than those within a single location less than 2 km apart ($F_{ST} = 0.04\sim 0.08$, $P < 0.01$). In contrast, a third site with a less impacted nearshore site (i.e., less pronounced environmental gradient) showed no significant structure from the offshore comparison. Our results show much stronger support for environment than distance separating these populations. Our finding suggests that ecological boundaries from human impacts may play a role in forming genetic structure in the coastal environment, and that genetic divergence in the absence of geographical barriers to gene flow might be explained by selective pressure across contrasting habitats.

Subjects Conservation Biology, Ecology, Genetics, Marine Biology

Keywords Coral reefs, Anthropogenic impacts, Population genetics, Hawaii, Lobe coral, Isolation by distance, Isolation by environment, Local adaptation, Gene flow

INTRODUCTION

Coral reefs are centers of marine biodiversity and productivity that provide a variety of ecosystem services of substantial cultural and economic value to humankind, yet coral reefs worldwide are under serious threat as a result of human activities (Hughes *et al.*, 2010; Graham, 2014). Average global coral cover has declined dramatically in the past 100 years due to a range of impacts such as sedimentation, pollution, overfishing, disease outbreaks and climate change (Hughes *et al.*, 2010; Richmond & Wolanski, 2011; Graham, 2014).

Submitted 18 November 2019

Accepted 12 January 2020

Published 18 February 2020

Corresponding authors

Kaho H. Tisthammer, ktist@sfsu.edu,

Ktisthammer@gmail.com

Robert J. Toonen,

rjtoonen@gmail.com,

toonen@hawaii.edu

Academic editor

Anastazia Banaszak

Additional Information and
Declarations can be found on
page 17

DOI 10.7717/peerj.8550

© Copyright

2020 Tisthammer et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Such effects are particularly pronounced in nearshore marine habitats, which are increasingly exposed to reduced water quality due to human activities ([Wenger et al., 2015](#)). Recent rapid coastal development, along with coastal industrial and recreational activities, have resulted in introducing sediments, nutrients and a variety of chemical pollutants to the nearshore environments ([Smith et al., 2008b](#); [Van Dam et al., 2011](#); [Wenger et al., 2015](#)). These local stressors often create a steep environmental gradient of water quality from nearshore toward offshore areas, and ‘signs of coral health impairment’ are usually detected along with the gradient (e.g., [Smith et al., 2008b](#); [Thompson et al., 2014](#); [Ennis et al., 2016](#)). Additionally, nearshore marine habitats are naturally exposed to higher fluctuations in temperature, pH and other environmental variables, creating contrasting environmental conditions relative to more stable offshore environments ([Gorospe & Karl, 2011](#); [Guadayol et al., 2014](#)). Some corals, however, continue to thrive in such nearshore ‘suboptimal’ habitats ([Morgan et al., 2016](#)), indicating that these individuals can withstand such stressors.

What impact does an ecological landscape with such a strong gradient have on the genetics of the organisms? ‘Isolation by distance’ (IBD, [Wright, 1943](#)) predicts that the degree of genetic differentiation increases with geographic distance due primarily to dispersal limits ([Slatkin, 1993](#); [Selkoe et al., 2016](#)). Isolation by distance is a neutral process in which dispersal limits gene flow and the scale over which genetic structure accumulates. ‘Isolation by environment’ (IBE, [Wang & Bradburd, 2014](#)) describes a pattern in which genetic differentiation increases with environmental differences, independent of geographic distance. Isolation by environment is a process that emphasizes the role of environmental heterogeneity and ecology in forming genetic structure, likely because of natural selection ([Orsini et al., 2013](#); [Wang & Bradburd, 2014](#)). Isolation by environment can be generated by different processes, including natural selection, sexual selection, reduced hybrid fitness, and biased dispersal; examples of the terms describing a specific case of IBE include ‘isolation by adaptation’ (IBA, [Nosil, Funk & Ortiz-Barrientos, 2009](#)), ‘isolation by colonization’ (IBC, [De Meester et al., 2002](#)), and ‘isolation by resistance’ (IBR, [McRae & Beier, 2007](#)). IBA and IBC emphasize the role of selection in forming genetic structure, and IBR describes correlation of genetic distance and resistance distance (i.e., friction to dispersal) ([McRae & Beier, 2007](#)). IBE along with related terms result in a pattern where genetic distance increases as ecological distance increases, but not with geographic distance for most loci ([Orsini et al., 2013](#); [Wang & Bradburd, 2014](#)). Theoretically, IBA, IBC and other processes will result in different distributions of genetic variation across landscapes ([Orsini et al., 2013](#)), though in reality, multiple processes almost always contribute to structuring genetic variation, and pinpointing the possible underlying processes may be difficult ([Selkoe et al., 2016](#)). For coastal marine ecosystems, often the distances between the impacted nearshore and un-impacted offshore sites are relatively small with no apparent dispersal barrier between adjacent sites for broadcast spawning species with pelagic larval development, providing an excellent opportunity to study IBE.

Several studies have shown coral genetic divergence occurring along ecological gradients. For example, Carlon and Budd ([Carlon & Budd, 2002](#)) described a pair of incipient species

in the coral *Favia fragum* associated with strong ecological gradients. The two types are largely restricted to alternate seagrass and adjacent coral reef habitats, but retain phenotypic distinction in a narrow zone of ecological overlap (Carlon & Budd, 2002). Subsequent work showed that the morphologies were heritable, and selection appeared to limit gene flow between the ecomorphs (Carlon et al., 2011). Carlon et al. (2011) postulated that divergent selection for “Tall” and “Short” ecomorphs of these inbred and brooding corals was driving the diversification of this coral via an ecological model of speciation (*sensu* Rundle & Nosil, 2005). Genetic divergence across nearshore and costal headland habitats has also been observed for a broadcast spawning coral *Porites lobata* in American Samoa (Barshis et al., 2010), and for brooding corals *Seriatopora hystrix* in Australia (Bongaerts et al., 2010) and *Porites astreoides* in the Florida Keys (Kenkel et al., 2013). Additionally, Gorospe & Karl (2015) found a significant genetic cline in *Pocillopora damicornis* along a depth gradient across a 40 m diameter patch reef in Hawai‘i.

Porites lobata (Dana, 1846), the study species, occurs over a wide geographic range in the tropical Pacific Ocean (Veron, 2000), and several studies have documented a pattern of isolation by distance across archipelagic (Polato et al., 2010) or broader scales (Baums et al., 2012), with little evidence of restricted gene flow among geographically proximate reefs at inter-island distances (but see Barshis et al., 2010). This massive coral is also known for its robustness; for example, *P. lobata* shows a high tolerance for sedimentation (Stafford-Smith, 1993) and bleaching (Levas et al., 2013), and a colony can recover from partial mortality due to tissues residing deep within the perforate skeleton, a phenomenon referred to as the ‘Phoenix effect’ (Roff et al., 2014). *Porites lobata* is one of the most dominant scleractinian coral species in Hawai‘i (Franklin, Jokiel & Donahue, 2013). Additionally, *P. lobata* shows high fidelity to a specific endo-symbiont, *Symbiodinium* Clade C15, which allows us to focus on responses of the host coral to environmental differences (LaJeunesse et al., 2004; Smith et al., 2008a; Barshis et al., 2010; Fabina et al., 2012).

Skeletal morphological differences for *P. lobata* appear to tell a similar story as for *F. fragum* (Forsman et al., 2015; Tisthammer & Richmond, 2018), though environmental gradients have not yet been tested in population genetic studies of this species in Hawai‘i. A growing number of studies indicate that IBE may be more ubiquitous in the sea than previously assumed. Therefore, we pose the question of whether there is reason to believe non-IBD patterns might be observed in contrasting habitats in Hawai‘i. In order to test our hypothesis, we implement a non-traditional population genetics sampling design that allows us to consider whether geography *vs.* environmental gradients explain overall patterns of genetic differentiation.

Maunalua Bay, Hawai‘i, O‘ahu, was selected as a study site due to the existence of a strong environmental gradient; large-scale urbanization in adjacent watersheds has caused severe deterioration in the health and extent of its nearshore coral reefs over the last century (Wolanski, Martinez & Richmond, 2009). Corals that survive in these affected nearshore areas are under chronic stress, and a previous survey showed significantly different cellular stress responses of individual colonies along the environmental gradient of pollutants and sedimentation from the inner bay toward

offshore (Wolanski, Martinez & Richmond, 2009; Richmond, 2009; Storlazzi et al., 2010). At the nearshore site of Maunaloa Bay, the suspended sediment concentration periodically exceeds several hundred mg/L, and the run-off water introduces toxicants such as benzo[a]pyrene, benzo[k]fluoranthene, phenanthrene and alpha-chlordane (Wolanski, Martinez & Richmond, 2009; Richmond, 2009; Storlazzi et al., 2010). The temperature, salinity and turbidity likewise all show higher ranges of values and fluctuations at the nearshore site than the offshore site (Storlazzi et al., 2010). The distance between the studied offshore and nearshore sites was less than 2 km across the extent of this gradient, and water movement in the areas suggests no dispersal barrier between the sites (Presto et al., 2012).

Wahikuli, West Maui, was selected as another study site due to the existence of a sharp environmental gradient that also runs from nearshore to offshore. Although less well characterized than O'ahu, the coral reefs off West Maui have a similar gradient of human impacts and experienced a dramatic decline in their coral cover from land-based sources of anthropogenic stressors over the last several decades (Rodgers et al., 2015). Substantial deterioration in the health of West Maui's coral reefs has lead Wahikuli and Honokōwai watersheds of West Maui to be designated as priority sites for conservation and management by the United States Coral Reef Task Force (USCRTF) and the State of Hawai'i (Williams et al., 2014). The Wahikuli study site is directly exposed to terrestrial run-off, due to its topography and current patterns (Storlazzi et al., 2006), causing high turbidity especially after heavy rains. Despite their proximity, the nearshore area at Wahikuli has markedly different water quality than offshore reefs roughly 300 m away (Fig. S2). In contrast, the nearshore area at the Honokōwai site is less affected by runoff, because it does not receive any direct stream discharge, resulting in consistently lower turbidity than the Wahikuli nearshore site (Vargas-Angel, 2017).

Porites lobata populations from Maunaloa Bay, O'ahu (hereafter O'ahu) and Wahikuli, Maui (hereafter Maui1) represent two locations with a strong environmental gradient, whereas Honokōwai, Maui (hereafter Maui2) represents a similar paired nearshore-offshore comparison site, but without a strong environmental gradient (Fig. 1). Nearshore and offshore sites at O'ahu and Maui1 sites are both characterized by highly contrasting environments in proximity, with dramatic differences in water quality and sedimentation loads from anthropogenic impacts. We undertook a genetic analysis of *P. lobata* across these sites to explore the possibility of isolation by environment. By comparing corals collected from heavily impacted nearshore environments to nearby congeners from more oceanic conditions, we sought to distinguish the roles played by ecology and anthropogenic impacts to the environment on the genetics of coral populations, in contrast to geographical distance limiting dispersal among similar habitats on adjacent islands. We predicted that the genetic structure of coral populations from areas with a strong environmental gradient would follow IBE, rather than IBD. At each location, we assessed the degrees of genetic differentiation and genetic diversity of *P. lobata* between adjacent strongly anthropogenically impacted 'higher-stress' nearshore and 'lower-stress' offshore sites, and compared them within and between locations to understand the effects of habitat types,




Figure 1 Maps of sampling locations. The Main Hawaiian Islands from which samples of *Porites lobata* were sampled. Paired sites at each location are designated as Nearshore (N) or Offshore (O) and geographic distances between the sampling sites are shown in gray.

Full-size [DOI: 10.7717/peerj.8550/fig-1](https://doi.org/10.7717/peerj.8550/fig-1)

anthropogenic impacts, and geographical distance on the genetic structure of reef building corals.

MATERIALS & METHODS

Species identification

Due to its high morphological variation, the genus *Porites* is notorious for its difficulties in distinguishing between its species (e.g., [Veron, 1995](#); [Veron, 2000](#); [Forsman et al., 2009](#); [Forsman et al., 2015](#)). Genetic delineation of some *Porites*, including *P. lobata*, has been challenging due to cryptic species and polymorphic or hybrid species complexes (e.g., [Veron & Pichon, 1982](#); [Forsman et al., 2009](#)). Although *Porites* corallites are small, irregular and can be highly variable, micro-skeletal (corallite) structures have been proposed to be important for species identification, therefore, we examined the corallites of all collected samples to confirm our taxonomic identifications ([Veron & Pichon, 1982](#); [Veron, 2000](#)) ([Fig. S3](#)). In Hawai'i, the only *Porites* species with a similar colony morphology to *P. lobata* is *Porites evermanni* (there are no records of *Porites lutea* in Hawai'i, although [Fenner \(2005\)](#) synonymized *P. evermanni* and *P. lutea*, they represent two distinct genetic clades ([Forsman et al., 2009](#)). *P. evermanni* is genetically distinct from *P. lobata* ([Forsman et al., 2009](#)), Clade V), and *P. lutea* has a distinct corallite skeletal morphology, compared to *P. lobata* ([Veron, 2000](#)).

Coral sampling

Small fragments (1 cm²) of *P. lobata* tissue samples were collected from live colonies between February 2013 to May 2017 at the following sampling sites in Hawai'i; (a) 'O'ahu'—nearshore ($n = 22$) and offshore ($n = 21$) sites at Maunaloa Bay, O'ahu (21.261~21.278°N, 157.711°W), (b) 'Maui1'—nearshore ($n = 21$) and offshore ($n = 23$) sites off the Hanakao'o Beach Park, West Maui (Wahikuli, 20.95°N, 156.68°W), and (c) Maui2—nearshore ($n = 23$), and offshore ($n = 20$) sites off the Honokōwai Beach Park, West Maui (Honokōwai, 20.90°N, 156.69°W) (Fig. 1, Fig. S1). Samples were taken from coral colonies at least two meters apart at each site, and after sampling, each coral colony was photographed and tagged to avoid resampling of the same colony. The collected tissue samples were either flash frozen in liquid nitrogen on shore and subsequently stored at -80, preserved in DMSO buffer (Gaither et al., 2011), or stored in 100% ethanol. Genomic DNA was extracted from each coral tissue sample using the Qiagen® DNeasy Blood & Tissue Kit. Coral samples were collected under the State of Hawai'i Division of Aquatic Resources, Special Activity Permit 2013-26, 2014-64, 2015-06, and 2017-16.

PCR

For the samples from O'ahu, the following three regions of coral host DNA were PCR-amplified: (1) ~400 bp coral mitochondrial putative control region (CR) with primers CRf and CO3r (Vollmer & Palumbi, 2002), (2) ~700 bp coral nuclear ITS1-5.8S-ITS2 region (ITS) with primers ITSZ1 and ITSZ2 (Forsman et al., 2009), and (3) ~1,500 bp coral nuclear histone region spanning H2A to H4 (H2) with novel primers zH2AH4f (5'-GTGTAAGTGGCTGCYGTCT-3') and zH4Fr (5'-GACAACCGAGAATGTCCGGT-3'). H2 was developed to create a genetic marker that allow direct sequencing of post PCR products to efficiently assess small-scale population genetic structure, because (1) the mitochondrial genome of *P. lobata* exhibits very little sequence variability (<0.02% polymorphic sites (Tisthammer et al., 2016)) due to its extremely slow evolutionary rate (Shearer et al., 2002), and (2) even though high polymorphism in ITS is a desirable trait, sequencing of ITS requires time-consuming cloning, and analyzing the multi-copy gene poses analytical challenges, as it deviates from a standard diploid model. H2 consists of partial coding region of H2A, approximately 1000 bp introns and part coding region of H4. After successful development of the H2 marker, Maui populations were analyzed with only H2 for the above reasons (see File S1 for more details).

H2 was amplified under the following conditions: 96 °C for 2 min (one cycle), followed by 34 cycles consisting of 96 °C for 20 s, 58.5 °C for 20 s, and 72 °C for 90 s, and a final extension at 72 °C for 5 min. H2 amplifications (25 µl) consisted of 0.5 µl of DNA template, 0.2 µl of GoTaq® DNA Polymerase (Promega, Madison, WI), 5 µl of GoTaq® Reaction Buffer, 1.6 µl of 50 mM MgCl₂, 2 µl of 10 mM dNTPmix, 1.6 µl of each 10 mM primer, and nuclease-free water to volume. For samples with multiple bands, approximately 1500-bp PCR products were extracted from agarose gels after electrophoresis and purified using the UltraClean® 15 DNA Purification Kit (MO BIO Laboratories, Carlsbad, CA)

according to the manufacturer's instruction. The rest of the PCR products were purified with UltraClean® PCR Clean-Up Kit (MO BIO Laboratories) and sequenced directly in both directions on the ABI 3730xl DNA Analyzer. Clone libraries were created for each amplified ITS region using the pGEM®-Easy Vector System (Promega). Positive inserts were verified by PCR using SP6 and T7 primers, and plasmids (2–5 per library) were treated with UltraClean® 6 Minute Mini Plasmid Prep Kit (MO BIO Laboratories) and sequenced on an ABI-3130XL Genetic Analyzer sequencer. For Maui samples, H2 was amplified and sequenced using the same method as described above.

Sequence analyses

Resulting DNA sequences were aligned using Geneious® 6.1.8 (Biomatters Ltd., Auckland, New Zealand). Polymorphic sites within H2 regions were identified using Geneious® (Find Heterozygotes option) and confirmed by eye. Middle sections, as well as both ends of H2 were then trimmed to 1,352 bp (for O'ahu sequences) or 1,221 bp (for combined O'ahu and Maui analysis) to minimize missing nucleotides among samples. H2 was phased using the program PHASE 2.1 (*Stephens, Smith & Donnelly, 2001*) and SeqPHASE (*Flot, 2010*). The analysis of molecular variance and other population genetic statistics were estimated in Arlequin 3.5 (*Excoffier & Lischer, 2010*) and TCS 1.21 (*Clement, Posada & Crandall, 2000*). The global AMOVA with a weighted average over loci with permutation tests was used as implemented in Arlequin 3.5, using pairwise difference as a distance computation method. For H2, both phased and non-phased sequences were run with AMOVA, which produced the same statistical results, and therefore only the results from the phased sequences are presented here. Up to five coral ITS sequences were successfully cloned and sequenced per colony, and the entire data set was used for calculation of population statistics, treating each cloned sequence as a haplotype. Attempts have been made to conduct genetic analysis using ITS by (a) treating each sequence as a haplotype (inclusivity), (b) making a consensus sequence per individual (consensus by plurality), or (c) using a hierarchical PERMANOVA (*Barshis et al., 2010*). In this study, we ran AMOVA using ITS by both (a) and (b) methods, which produced the same statistical outcome, and hence, the results from inclusivity (a) are presented in this paper. To address the unequal sample sizes (28 vs 44) between the sites in Maunalua Bay, the analysis was repeated after resampling to the equal sample size (28) for 10 times. All DNA sequences were inspected for the possibility of multi-sampled individuals, and all sampled colonies were considered as separate individuals (genets) since no two individuals from a single site shared the same haplotypes (H2). Mantel's test for isolation by distance was run on the samples in R (*R Core Team, 2014*) using pairwise genetic distance with 5000 bootstrap permutations. Rarefaction analysis was conducted in Analytic Rarefaction 2.1.1 (*Holland, 2009*). The network analysis was conducted in SplitsTree v.4.14.2, using UncorrectedP as a distance method, NeighborNet as a network computing method.

Table 1 Analysis of Molecular Variance (AMOVA) for *Porites lobata* collected from O'ahu (Maunalua Bay).

	Source of Variation	Variance components	% Variance	F_{ST}
ITS ($n = 70$)	Between populations	2.27	19.18	0.19^{***}
	Within populations	9.56	80.82	
H2 ($n = 43$)	Between populations	0.29	7.15	0.072^{***}
	Within populations	1.30	31.88	
	Within individuals	2.49	60.96	
CR ($n = 20$)	Between populations	0.034	8.49	0.086
	Within populations	0.370	91.5	($P = 0.15$)

Notes. Populations here refer simply to individuals sampled within the same sampling location, with sample size in parentheses below the marker. % Variance refers to the proportion of genetic variation explained by each comparison, and bold numbers with *** are significant at $P < 0.001$.

RESULTS

Nearshore vs. offshore comparison of genetic structure and diversity of *P. lobata* populations O'ahu (Maunalua Bay)

For O'ahu *P. lobata* populations, the degree of genetic differentiation was estimated using analysis of molecular variance (AMOVA (Excoffier & Lischer, 2010)) between the nearshore and offshore sites using three genetic markers; CR, ITS, and novel H2 developed for this study. The AMOVA results for both nuclear markers revealed clear genetic differentiation between the two sites (ITS, $F_{ST} = 0.19$, $P < 0.001$; H2, $F_{ST} = 0.072$, $P < 0.001$) (Table 1). The mitochondrial marker (CR) did not detect significant differentiation ($F_{ST} = 0.086$, $P = 0.15$), which was not surprising due to its extremely low variability in corals and cnidarians in general (Shearer et al., 2002). The numbers of shared haplotypes (alleles) between the nearshore and offshore O'ahu populations were also low; out of 37 ITS haplotypes identified from the 70 total sequences, only three (8.1%) were shared between the sites. For H2, there were 54 unique haplotypes out of 86 total phased sequences, and only 5 sequences (9.3%) were shared between the sites (Table 2). The pattern of genetic structure was visualized using network analysis, which revealed sequences clustering into three major groups in both ITS and H2 markers, which consisted of one cluster dominated by the nearshore individuals, the second one dominated by the offshore individuals, and the last group with approximately mixed origins (Fig. 2). For CR, three haplotypes were identified from 27 sequences, all of which were present at both sites. Interestingly, the most common haplotype was the most dominant one at the nearshore site, while the second common haplotype was the dominant haplotype at the offshore site, though the AMOVA, as well as the exact test results were not significant ($P = 0.15$ and 0.08 respectively) (Fig. 3).

The pattern of genetic diversity also differed between the nearshore and offshore populations. The degree of genetic diversity was higher at the offshore site; percent private alleles (pA), percent polymorphic sites (poly), and nucleotide diversity level (π) were almost twice as high in the offshore population as in the nearshore one based on ITS (Table 2). Standardizing sample size by random resampling confirmed that this was not

Table 2 Population genetic statistics of *Porites lobata* from O'ahu (Maunalua Bay).

Sites	ITS (707 bp)										
	<i>n</i>	A	pA	poly	D_A	D_P	<i>i</i>	π	$\theta\pi$	θ_s	
O'ahu Nearshore	28	13 (46%)	10 (36%)	45 (6.4%)	1.0 ± 0.0095	0.259 ± 0.182	31	0.0167 ± 0.009	11.64 ± 5.44	3.60 ± 1.45	
O'ahu Offshore	42	27 (64%)	24 (57%)	70 (10%)	1.0 ± 0.0052	0.343 ± 0.192	50	0.0340 ± 0.017	24.03 ± 10.78	6.04 ± 2.06	
O'ahu Offshore*	(28)	21.7 (78%)	19.3 (69%)	65.2 (9.2%)			48.5	0.0337 ± 0.017	23.7 ± 11.96	5.37 ± 2.00	

Sites	H2 (1,352 bp)										
	<i>n</i>	A	pA	poly	H_o	H_e (D_A)	D_P	hom	π	$\theta\pi$	θ_s
O'ahu Nearshore	22 (44)	28 (64%)	23 (52%)	27 (2.0%)	0.773	0.965	0.120 ± 0.151	5 (23%)	0.00553 ± 0.003	7.483 ± 3.96	6.207 ± 2.09
O'ahu Offshore	21 (42)	31 (74%)	26 (62%)	27 (2.0%)	0.810	0.977	0.162 ± 0.179	4 (19%)	0.00558 ± 0.003	7.554 ± 4.00	6.275 ± 2.12

Sites	CR (366 bp)										
	<i>n</i>	A	pA	poly	D_A	D_P	π	$\theta\pi$	θ_s		
O'ahu Nearshore	13	3 (23%)	0 (0%)	2 (0.5%)	1.0 ± 0.0302	0.282 ± 0.000	0.00154 ± 0.0015	0.5641 ± 0.551	0.6445 ± 0.485		
O'ahu Offshore	14	3 (21%)	0 (0%)	2 (0.5%)	0.45 ± 0.0270	0.451 ± 0.124	0.0056 ± 0.003	0.9011 ± 0.747	0.6289 ± 0.474		

Notes.

Sample size (*n*, for H2, the number in parentheses represents the number of phased sequences), number of haplotypes (A), number of private haplotypes (pA), number of polymorphic sites (poly), mean overall gene diversity ($D_A \pm SD$), mean gene diversity for polymorphic sites only ($D_P \pm SD$), observed heterozygosity (H_o), expected heterozygosity (H_e), number of indels (*i*), number of homozygous individuals (hom), nucleotide diversity ($\pi \pm SD$), theta estimator 1 ($\theta\pi$: expected heterozygosity at a nucleotide position estimated from the mean π), theta estimator 2 (Watterson estimator, θ_s).

*Standardized values to the minimum sample size of 28.

an artifact of a larger sample size of the offshore population (Table 2). Rarefaction analysis of ITS sequences also confirmed that allelic richness of the offshore population (Richness = 21.6 ± 2.1 at $n = 28$, 95% CI [18.8–24.5]) was clearly higher than that of the nearshore population (Richness = 13) (Fig. 4). The level of genetic diversity in H2 also appeared slightly higher in the offshore populations; the number of haplotypes (A), the number of private allele (pA), the heterozygosity level (H_o), the number of heterozygous individuals, and mean gene diversity (D_A , D_P) all had marginally higher values in the offshore samples, though the differences were small (Table 2). In both nuclear markers, $\theta\pi$ (the expected heterozygosity estimated from the average nucleotide diversity) was higher than θ_s (the theta estimated from the number of segregating sites).

Maui

At the two study locations on the island of Maui, patterns of genetic structure of *P. lobata* populations between the nearshore and offshore sites were analyzed using the novel H2 marker (see Material and Methods). At Maui1, where a strong environmental gradient exists, significant genetic differentiation was detected ($F_{ST} = 0.0415$, $P = 0.0308$), but the level of genetic diversity was comparable between the two sites (Table 3). At Maui2,




Figure 2 Allele networks for *Porites lobata* sampled from O'ahu (Maunaloa Bay). NeighborNet phylogenetic networks generated by SplitsTree, based on (A) ITS and (B) H2 color-coded by Nearshore (yellow) and Offshore (blue) sampling locations as shown in Fig. 1. Pie charts represent the proportion of sequences within each cluster.

Full-size [DOI: 10.7717/peerj.8550/fig-2](https://doi.org/10.7717/peerj.8550/fig-2)




Figure 3 Mitochondrial haplotype network for *Porites lobata* sampled from O'ahu (Maunaloa Bay). Haplotype network based on mitochondrial putative control region (CR) color-coded by Nearshore and Offshore sampling locations as shown in Fig. 1.

Full-size [DOI: 10.7717/peerj.8550/fig-3](https://doi.org/10.7717/peerj.8550/fig-3)

which had much less contrasting environmental conditions between the nearshore and offshore sites, the AMOVA results found no significant genetic differentiation between the nearshore and offshore sites ($F_{ST} = 0.0019$, $P = 0.991$). The level of genetic diversity at Maui2 appeared slightly higher in the nearshore population, which had higher numbers of haplotypes (A), polymorphic sites (poly), and heterozygous individuals (Table 3). The theta estimators of Maui2 also showed a different pattern from the O'ahu and Maui1 populations, with higher values of θ_s than those of θ_π at both nearshore and offshore sites.




Figure 4 Rarefaction curve of allelic richness of ITS sequences from *Porites lobata* sampled from nearshore and offshore populations on O'ahu (Maunalua Bay). Allelic richness from individuals collected at Nearshore and Offshore sampling sites of O'ahu as shown in Fig. 1. The gray bar indicates the 95% confidence interval of the allelic richness estimation for the offshore population at equivalent sample size of $n = 28$.

Full-size DOI: 10.7717/peerj.8550/fig-4

Table 3 Population genetic statistics of *Porites lobata* from Maui1 and Maui2 sites.

	H2 (1,221 bp)									
	<i>n</i>	A	pA	poly	hom	Ho	He	π	$\theta\pi$	θ_s
Maui1	18	31	15	27	3	0.833	0.987	0.00596 ± 0.0032	7.271 ± 3.87	6.511 ± 2.25
Nearshore	(36)	(86%)	(83%)	(2.2%)	(17%)					
Maui1	22	35	18	31	4	0.818	0.986	0.00590 ± 0.0031	7.209 ± 3.82	7.126 ± 2.34
Offshore	(44)	(80%)	(82%)	(2.5%)	(18%)					
Maui2	23	38	43	65	3	0.870	0.988	0.00614 ± 0.0035	7.494 ± 3.96	14.790 ± 4.48
Nearshore	(46)	(86%)	(93%)	(5.3%)	(13%)					
Maui2	20	30	37	33	4	0.800	0.967	0.00481 ± 0.0026	5.878 ± 3.19	7.758 ± 2.57
Offshore	(40)	(75%)	(93%)	(2.7%)	(20%)					

Notes.

Based on 1,319 bp of H2 sequence data. Sample size (n , represents the number of phased sequences), number of haplotypes (A), number of private haplotypes (pA), number of polymorphic sites (poly), mean overall gene diversity ($D_A \pm SD$), mean gene diversity for polymorphic sites only ($D_P \pm SD$), observed heterozygosity (H_O), expected heterozygosity (He), number of indels (i), number of homozygous individuals (hom), nucleotide diversity ($\pi \pm SD$), theta estimator 1 ($\theta\pi$: expected heterozygosity at a nucleotide position estimated from the mean π), theta estimator 2 (Watterson estimator, θ_s). * Standardized values to the minimum sample size of 28.

O'ahu vs. Maui

Inter-island genetic structure, as well as comparison of nearshore and offshore populations were conducted using H2 marker. The hierarchical AMOVA did not detect significant structure between the O'ahu and Maui populations ($F_{CT} = 0.007$, $P = 0.27$), but the two Maui locations showed significant differentiation ($F_{SC} = 0.063$, $P = 0.000$) based on H2 (Table S2A). The patterns of genetic diversity suggested an overall lower variability in the O'ahu population; the numbers of haplotypes (A), polymorphic sites (poly), and

Table 4 Population genetic statistics of *Porites lobata* from the islands of Oahu and Maui.

	H2 (1,221 bp)									
	<i>n</i>	A	pA	poly	hom	Ho	He	π	$\theta\pi$	θ_s
Maui1 (pooled)	40 (80)	63 (79%)	76 (95%)	38 (3.1%)	7 (15.9%)	0.825	0.986	0.0060 ± 0.0032	7.385 ± 3.87	7.672 ± 2.26
Maui2 (pooled)	43 (86)	68 (79%)	82 (95%)	82 (6.7%)	7 (16.2%)	0.837	0.983	0.00571 ± 0.0030	6.972 ± 3.67	16.316 ± 4.38
Maui (pooled)	83 (166)	126 (76%)	159 (81.5%)	96 (7.9%)	14 (16.1%)	0.831	0.987	0.00604 ± 0.0031	7.380 ± 3.84	16.355 ± 3.96
O'ahu (pooled)	43 (86)	54 (62.7%)	47 (87%)	35 (2.9%)	9 (20.9%)	0.791	0.974	0.00643 ± 0.0033	7.844 ± 4.08	6.964 ± 2.06

Notes.

Based on 1,221 bp of H2 sequence data. Sample size (*n*, represents the number of phased sequences), number of haplotypes (A), number of private haplotypes (pA), number of polymorphic sites (poly), mean overall gene diversity ($D_A \pm SD$), mean gene diversity for polymorphic sites only ($D_P \pm SD$), observed heterozygosity (H_O), expected heterozygosity (He), number of indels (i), number of homozygous individuals (hom), nucleotide diversity ($\pi \pm SD$), theta estimator 1 ($\theta\pi$: expected heterozygosity at a nucleotide position estimated from the mean π), theta estimator 2 (Watterson estimator, θ_s). * Standardized values to the minimum sample size of 28.

heterozygous individuals were all smaller on O'ahu, although nucleotide diversity (π) levels were relatively similar between O'ahu and Maui (Table 4).

Pairwise F_{ST} comparisons between all combinations revealed that the nearshore populations from O'ahu and Maui1 with a high level of environmental stress were genetically closer to each other than to their respective, nearby offshore populations, and similarly the offshore populations from O'ahu and Maui1 were genetically closer to each other than to their respective nearshore populations (Table 5, O'ahu and Maui1 populations). Assessing by habitat types, the nearshore and offshore populations of O'ahu and Maui1 also exhibited significant genetic differentiation ($F_{ST} = 0.065$, $P = 0.000$, Table S2B). The results also revealed that Maui2 corals, which showed no significant structure between the nearshore and offshore sites, turned out to be rather genetically unique compared to the rest of the populations. However, the F_{ST} values indicated that the Maui2 nearshore population was genetically closer to other offshore populations ($F_{ST} = 0.027$ – 0.035) than to other nearshore populations of O'ahu and Maui1 ($F_{ST} = 0.14$ – 0.18), suggesting collectively that Maui2 corals at both sites were genetically closer to the offshore populations (Table 5).

Genetic structure of *P. lobata* across islands was also visualized using network analysis, which revealed three major clusters of H2 sequences, similar to the results from the O'ahu populations (Fig. 5). Grouping by habitat-based genetic groups, Cluster 1 was dominated by the offshore type (including Maui2-nearshore) (88%), Cluster 2 was dominated by nearshore individuals (74%), and Cluster 3 had approximately same proportion of nearshore and offshore types, depicting separation of offshore and nearshore individuals, especially for O'ahu and Maui1 populations. No clear pattern was observed based on geographic locations (Fig. S4).

DISCUSSION

Previous work on *P. lobata* at the Archipelagic scale (Polato et al., 2010) found IBD, although habitat-level variation was not examined. At a much smaller spatial scale

Table 5 Pairwise F_{ST} values for *Porites lobata* sampled from Oahu and Maui.

	O'ahu N	O'ahu O	Maui1 N	Maui1 O	Maui2 N	Maui2 O
O'ahu N	–	0.000	0.324	0.000	0.000	0.000
O'ahu O	0.079^{***}	–	0.008	0.666	0.012	0.046
Maui1 N	0.001	0.055^{**}	–	0.004	0.000	0.000
Maui1 O	0.077^{***}	–0.007	0.052^{**}	–	0.015	0.014
Maui2 N	0.177^{***}	0.035[†]	0.139^{***}	0.027[†]	–	0.072
Maui2 O	0.205^{***}	0.026[†]	0.170^{***}	0.030[†]	0.013	–

Notes.

Pairwise F_{ST} values were estimated using AMOVA in Arlequin with 5,000 permutations. Below diagonal = F_{ST} values, Above diagonal = P values.

Bold values are significant, and asterisks represent significance values (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$).

N, Nearshore; O, Offshore.




Figure 5 Diagrams of neighbor-net tree networks for O'ahu and Maui *P. lobata* populations based on unphased H2 sequences. Genetic structure is stronger across human-impacted habitats than among islands in the coral *Porites lobata* NeighborNet phylogenetic networks generated by SplitsTree, color-coded by habitat-based genetic clusters: Blue represents the offshore group (including Maui2-nearshore population), and yellow represents the two genetically-close nearshore populations of O'ahu and Maui1. The pie charts show the proportion of sequences present in each group. The gray numbers in () represent the proportion of Maui2 nearshore population.

Full-size DOI: 10.7717/peerj.8550/fig-5

considered here, with explicit sampling of contrasting habitats from across this strong environmental gradient, the pattern of genetic structure we observed for *P. lobata* in Hawai'i did not follow IBD (Mantel Test, $r = -0.091$, $P = 0.54$). Instead, a pattern

that resembles IBE was revealed, with a correlation between habitat types irrespective of geographic distance; the pairwise F_{ST} values revealed that offshore individuals from two separate islands (>100 km) were genetically closer to each other than to their geographically closest nearshore individuals (0.3–2 km), and nearshore individuals from two islands were also typically genetically closer to each other than to corals from adjacent offshore sites (Table 5, Fig. 5). Other studies that examined genetic differentiation on multiple scales show that patterns of IBD can be scale dependent. For example, Gorospe and Karl found distance (2013) and depth (2015) dependent patterns of coral genetic relatedness on a single patch reef, but this pattern was no longer detectable for analyses on an inter-reef scale (Gorospe & Karl, 2013). Furthermore, because of intra-reef patterns of genetic diversity and clonality, it is possible that sampling design may bias population-level patterns of genetic diversity (Gorospe, Donahue & Karl, 2015). However, we do not think this potential bias presents a problem in our system, since *P. lobata* is a broadcasting spawning species and shows little clonality (Boulay et al., 2012; Boulay et al., 2013; Schweinsberg, Tollrian & Lampert, 2016), particularly in Hawai'i (Polato et al., 2010).

Similar patterns of small-scale genetic structure have been observed in several reef building corals (Barshis et al., 2010; Bongaerts et al., 2010; Carlon & Lippé, 2011). In the case of *Favia fragum*, the “Tall” ecomorph is a seagrass specialist with morphological adaptations to minimize sediment impacts, whereas the “Short” ecomorph shows morphological specializations for coral reef habitats that decrease its fitness in seagrass beds (Carlon & Budd, 2002). These traits are highly heritable and divergent selection appears to be driving reproductive isolation among these morphs and ongoing diversification in these incipient species (Carlon & Lippé, 2011). Here, we find a similar pattern of repeated genetic differentiation across strong ecological gradients driven by anthropogenic impacts (between adjacent sites 0.3–2 km apart), but little evidence of differentiation among similar habitats more than 100 km distant. Moreover, corals from a control site that lacks such a strong environmental gradient (Maui2) did not show the significant genetic structure documented at the other sites, which further suggests the potential role of environment in forming the observed genetic patterns (IBE).

Additionally, for the O'ahu populations, differences in microskeletal morphology have been reported between the nearshore and offshore sites (Tisthammer & Richmond, 2018), similar to the correlation seen between genetic distance and microskeletal morphology across broader geographic regions in *P. lobata* (Forsman et al., 2015). The most noticeable difference was the height of pali (inner vertical skeletal structure that usually exists in a set of eight in *P. lobata*) within a corallite (the structure associated with individual polyps); nearshore corals had taller and more pronounced pali than the offshore ones. Because exact functions and heritability of the traits are unknown, whether the observed morphological differences are due to divergent selection cannot be answered at this point. However, the study suggested that the differences might be due to potential beneficial roles played by larger pali in shedding sediments in turbid water (Tisthammer & Richmond, 2018), similar to the case of the Caribbean coral *F. fragum* (Carlon & Budd, 2002; Carlon & Lippé, 2011; Carlon et al., 2011). Correlation between the morphological and genetic distances reported

in the study is consistent with the idea of the likely role of environment in observed divergence.

Although we cannot rule out the possibility of prevailing oceanographic currents or barriers playing a role in producing our observed pattern of genetic patchiness, detailed studies on water movement suggests little dispersal barrier between the nearshore and offshore sites. In the bay at the O`ahu site, surface currents primarily flow west due to the prevailing trade-winds (offshore to nearshore). The below surface current movement seems to be more complex, and is generally towards the east (nearshore to offshore) with the presence of small eddies, at least during the summer ([Presto et al., 2012](#)). Eddies would increase the larval retention time in the bay during the summer spawning season, especially for *Porites* species that produce neutrally buoyant gametes ([Hunter, 1988](#)). For Maui sites, the detailed water movement around the study areas revealed the presence of shear zones between the shallower (<15 m) nearshore and the deeper portions of the reef due to opposing flow directions during the coral reproductive season (summer), suggesting local retention of larvae ([Storlazzi et al., 2006](#); [Storlazzi & Field, 2008](#)). While these shear zones could contribute as a gene flow barrier between the shallower and deeper areas, our offshore sites were located within the nearshore southward-flow dominated area, based on their depths and distances from the shore, thereby eliminating the probability of existence of strong oceanographic barriers between the nearshore and offshore study sites.

It is particularly interesting to find not only this pattern of clear genetic partitioning within a bay is repeated on a neighbor island, but also the genetic similarity exists between sites with similar histories of anthropogenic stress from the two separate islands, which points to the possibility of local processes acting to homogenize these geographically-separated sites. The nearshore sites at O`ahu and Maui1 have experienced deteriorating water and substrate quality due to terrestrial runoff from urbanization of adjacent watersheds over the past century. This repeated pattern implies a possibility of similar underlying processes at both locations. We suspect that the environmental decline has likely limited new recruitment to the affected nearshore sites ([Puritz & Toonen, 2011](#)), and may have placed the populations under local selection, similar to what was seen in the Caribbean coral *F. fragum* ([Carlon & Budd, 2002](#); [Carlon & Lippé, 2011](#)). The genetic markers used here are unlikely to be the direct targets of selection, and therefore, the observed genetic-environment association may result from linkage or be attributed to coupling of endogenous (intrinsic) and exogenous barriers, as theorized by [Bierne et al. \(2011\)](#). According to [Bierne et al. \(2011\)](#), intrinsic barriers that are independent of environment often play an important role in forming genetic-environmental associations through coupling with exogenous factors, rather than ecologically based divergent selection alone. Although our understanding on how selection affects population differentiation at neutral markers is limited at this point ([Bierne et al., 2011](#)), the coupling hypothesis may explain parallel genetic differentiation across independent putatively neutral markers observed in our study. Furthermore, there appears to be reduced genetic diversity in the nearshore habitats ([Table 2](#), [Table S1](#)), which may represent a subset of the standing genetic variation of the larger population, thereby representing a limited number of individuals capable of surviving in the nearshore habitats. Regardless of the precise

underlying mechanism, the role of environment in shaping the observed genetic divergence appears more important than geographic distance in this case.

Additional work is also needed to determine the specific environmental drivers likely to result in selection across these environmental gradients that generate the observed pattern. As discussed earlier, there are many factors, both natural and anthropogenic, that contrast between nearshore and offshore environments (e.g., salinity, irradiance, UV exposure, temperature, pH, wave exposure, nutrients, and biological community). Also, fine-scale water movements that the oceanographic-scale studies cannot capture (isolation by resistance, [Thomas et al., 2015](#)) may be contributing to physical barriers to reproduction. Any of these factors could contribute to create genetic partitioning between nearshore and offshore sites. For example, a comparable pattern of genetic structure has been observed across a particularly strong temperature gradient between the back-reef and forereef *P. lobata* populations in the areas with negligible terrestrial runoff or pollution in American Samoa ([Barshis et al., 2010](#)). However, in Hawai'i we did not see such differentiation between nearshore and offshore sites at our less-impacted reference (Maui2), which raises a question of a potential role of other local abiotic or biotic factors affecting the observed pattern. It will be interesting to continue to observe whether this differentiation is transient and of little evolutionary importance, or whether it progresses towards incipient speciation, as it appears to have done in the Caribbean coral *F. fragum* ([Carlon & Budd, 2002](#); [Carlon & Lippé, 2011](#)). Indeed, our finding may shed light on the common pattern of 'chaotic genetic patchiness' ([Johnson & Black, 1982](#)) so commonly reported among population genetic studies of marine organisms, in which geographically proximate populations show greater genetic differentiation than those from distant sites (e.g., [Johnson & Black, 1984](#); [Hogan, Thiessen & Heath, 2010](#); [Toonen & Grosberg, 2011](#); [Eldon et al., 2016](#)).

The ecological diversification of reef building corals over a small spatial scale, despite ongoing gene flow, also provides a rare example of genetic divergence in the absence of spatial barriers to gene flow, indicating that divergent natural selection can act as an evolutionary driver of reproductive isolation ([Carlon & Budd, 2002](#); [Bongaerts et al., 2010](#); [Bongaerts et al., 2011](#); [Carlon et al., 2011](#)). Our results may extend another example of *P. lobata* in Hawai'i to these findings, which showed potential occurrence of similar diversification process across steep environmental gradients driven. This may represent the initial stages of adaptive diversification, as seen in other marine species from the Hawaiian Archipelago (e.g., limpets, [Bird et al., 2011](#); [Bird, 2011](#)). There is clearly some genetic connectivity among adjacent islands, congruent to previous studies ([Polato et al., 2010](#)), and hence the observed differentiation across these steep ecological gradients in spite of high dispersal potential ([Storlazzi & Field, 2008](#); [Presto et al., 2012](#)) may be the early phases of speciation with gene flow ([Feder, Egan & Nosil, 2012](#)). Whether this initial stage of divergence among habitats is transient or has the potential to progress to later stages remains to be seen, but our results and others ([Bird et al., 2011](#); [Bird, 2011](#)) indicate that this initial stage can be realized even in a broadcasting species with high dispersal potential. Together, these results add to the growing evidence that the initial phase of speciation is possible without geographic isolation, and lend support to the hypothesis

that ecological speciation (*sensu Bowen et al., 2013*) may be more common in the sea than believed previously.

Finally, the potential reduction in intraspecific genotypic diversity within the populations of corals under higher levels of environmental stressors can be a warning for the future of reefs. Focusing on the usual metrics of coral reef health and resilience, specifically coral cover and species diversity, misses the very concerning loss of genotypic diversity essential to population resilience in a changing world. The importance of molecular data is apparent in tracking the invisible changes that can lead to local extinctions, and hence, such tools need to be further refined and more broadly applied to the helping ensure the persistence of coral reefs as a legacy for the future.

CONCLUSIONS

Our results show that the broadly distributed broadcast spawning coral *P. lobata* exhibits very fine-scale (0.3~2 km) genetic structure, and environmental drivers across habitat types appear to have a stronger effect in forming such genetic structure (IBE) than geographic distances (IBD) in coastal areas that are heavily affected by anthropogenic stressors. Genetic similarity found between O`ahu and Maui1 nearshore populations suggest that the observed genetic structure may be driven by similar underlying evolutionary mechanisms, including adaptation to environmentally driven factors. Our results also highlight the importance of thorough sampling among habitats at small scales; we could easily overlook such important local genetic differences, and may mistakenly conclude that populations are uniform across the landscape without thorough sampling. Although our samples are from limited locations, these results demonstrate that understanding small-scale genetic variation and diversity can provide important information on the ecological basis of genetic diversity and differentiation, which must be understood to effectively implement future coral reef conservation efforts.

ACKNOWLEDGEMENTS

We thank the following people for assistance in collecting corals: M Stiber, V Sindorf, F Seneca, J Murphy, J Martinez, A Lyman, K Richmond, V Hölzer, N Spies, and A Irvine. Special thanks to T Oliver for his support and advice. ZHF would like to thank the Seaver Institute for support.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

Financial support came from the National Fish and Wildlife Foundation (Grant Number 34413), the Hawaii Department of Health (MOA 13-502), and the National Oceanic and Atmospheric Administration, Coral Reef Ecosystem Studies Program (NA17NMF4520144) to Robert H. Richmond. Zac H. Forsman was supported by awards from the Seaver Institute and NSF-OA#1416889 to Robert J. Toonen. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

National Fish and Wildlife Foundation: Grant Number 34413.

Hawaii Department of Health: MOA 13-502.

National Oceanic and Atmospheric Administration, Coral Reef Ecosystem Studies Program: NA17NMF4520144.

Seaver Institute and NSF-OA#1416889.

Competing Interests

Robert J. Toonen is an Academic Editor and Section Editor for Aquatic Biology for PeerJ.

Author Contributions

- Kaho H. Tisthammer conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Zac H. Forsman conceived and designed the experiments, performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Robert J. Toonen analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Robert H. Richmond conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Coral samples were collected under the State of Hawaii Division of Aquatic Resources, Special Activity Permits #2013-26, 2014-64, 2015-06, and 2017-16.

Data Availability

The following information was supplied regarding data availability:

All raw sequence data are available at GenBank:

ITS: [KY493091–KY493160](#); H2: [KY502280–KY502376](#) and [MF629151–MF629152](#); CR: [KY502373](#), [KY502374](#) and [KY502375](#).

The phased H2 sequences used in the analysis are available on Figshare: Toonen, Rob (2019): Phased_H2_sequence_data.docx. figshare. Dataset. <https://doi.org/10.6084/m9.figshare.10315430.v1>.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8550#supplemental-information>.

REFERENCES

- Barshis DJ, Stillman JH, Gates RD, Toonen RJ, Smith LW, Birkeland C. 2010. Protein expression and genetic structure of the coral *Porites lobata* in an environmentally

- extreme Samoan back reef: does host genotype limit phenotypic plasticity? *Molecular Ecology* **19**:140297–140297 DOI [10.1111/j.1365-294X.2010.04574.x](https://doi.org/10.1111/j.1365-294X.2010.04574.x).
- Baums IB, Boulay JN, Polato NR, Hellberg ME. 2012.** No gene flow across the Eastern Pacific Barrier in the reef-building coral *Porites lobata*. *Molecular Ecology* **21**:5418–5433 DOI [10.1111/j.1365-294X.2012.05733.x](https://doi.org/10.1111/j.1365-294X.2012.05733.x).
- Bierne N, Welch J, Loire E, Bonhomme F, David P. 2011.** The coupling hypothesis: why genome scans may fail to map local adaptation genes. *Molecular Ecology* **20**:2044–2072 DOI [10.1111/j.1365-294X.2011.05080.x](https://doi.org/10.1111/j.1365-294X.2011.05080.x).
- Bird CE. 2011.** Morphological and behavioral evidence for adaptive diversification of sympatric Hawaiian Limpets (*Cellana* spp.). *American Zoologist* **51**:466–473 DOI [10.1093/icb/icr050](https://doi.org/10.1093/icb/icr050).
- Bird CE, Holland BS, bowen BW, Toonen RJ. 2011.** Diversification of sympatric broadcast-spawning limpets (*Cellana* spp.) within the Hawaiian archipelago. *Molecular Ecology* **20**:2128–2141 DOI [10.1111/j.1365-294X.2011.05081.x](https://doi.org/10.1111/j.1365-294X.2011.05081.x).
- Bongaerts P, Riginos C, Hay KB, Van Oppen MJ, Hoegh-Guldberg O, Dove S. 2011.** Adaptive divergence in a scleractinian coral: physiological adaptation of *Seriatopora hystrix* to shallow and deep reef habitats. *BMC Evolutionary Biology* **11**:303 DOI [10.1111/j.1365-294X.2011.05081.x](https://doi.org/10.1111/j.1365-294X.2011.05081.x).
- Bongaerts P, Riginos C, Ridgway T, Sampayo EM, Van Oppen MJH, Englebert N, Vermeulen F, Hoegh-Guldberg O. 2010.** Genetic divergence across habitats in the widespread Coral *Seriatopora hystrix* and its associated *Symbiodinium*. *PLOS ONE* **5**:e10871 DOI [10.1371/journal.pone.0010871.t001](https://doi.org/10.1371/journal.pone.0010871.t001).
- Boulay JN, Cortes J, Nivia-Ruiz J, Baums IB. 2012.** High genotypic diversity of the reef-building coral *Porites lobata* (Scleractinia: Poritidae) in Isla del Coco National Park, Costa Rica. *Revista de Biología Tropical* **60**:279–292.
- Boulay JN, Hellberg ME, Cortes J, Baums IB. 2013.** Unrecognized coral species diversity masks differences in functional ecology. *Proceedings of the Royal Society B: Biological Sciences* **281**:20131580–20131580 DOI [10.1111/j.1365-294X.2005.02553.x](https://doi.org/10.1111/j.1365-294X.2005.02553.x).
- Bowen BW, Rocha LA, Toonen RJ, Karl SA, Laboratory TT. 2013.** The origins of tropical marine biodiversity. *Trends in Ecology & Evolution* **28**:359–366 DOI [10.1016/j.tree.2013.01.018](https://doi.org/10.1016/j.tree.2013.01.018).
- Carlson DB, Budd AF. 2002.** Incipient speciation across a depth gradient in a scleractinian coral? *Evolution* **56**:2227–2242 DOI [10.1111/j.0014-3820.2002.tb00147.x](https://doi.org/10.1111/j.0014-3820.2002.tb00147.x).
- Carlson DB, Budd AF, Lippé C, Andrew RL. 2011.** The quantitative genetics of incipient speciation: heritability and genetic correlations of skeletal traits in populations of diverging fava fragum ecomorphs. *Evolution* **65**:3428–3447 DOI [10.1111/j.1558-5646.2011.01389.x](https://doi.org/10.1111/j.1558-5646.2011.01389.x).
- Carlson DB, Lippé C. 2011.** Estimation of mating systems in Short and Tall ecomorphs of the coral *Favia fragum*. *Molecular Ecology* **20**:812–828 DOI [10.1111/j.1365-294X.2010.04983.x](https://doi.org/10.1111/j.1365-294X.2010.04983.x).
- Clement M, Posada D, Crandall KA. 2000.** TCS: a computer program to estimate gene genealogies. *Molecular Ecology* **9**:1657–1659 DOI [10.1046/j.1365-294x.2000.01020.x](https://doi.org/10.1046/j.1365-294x.2000.01020.x).

- De Meester L, Gómez A, Okamura B, Schwenk K. 2002.** The Monopolization Hypothesis and the dispersal–gene flow paradox in aquatic organisms. *Acta Oecologica* **23**:121–135 DOI [10.1016/S1146-609X\(02\)01145-1](https://doi.org/10.1016/S1146-609X(02)01145-1).
- Eldon B, Riquet F, Yearsley J, Jollivet D, Broquet T. 2016.** Current hypotheses to explain genetic chaos under the sea. *Current zoology* **62**:551–566 DOI [10.1093/cz/zow094](https://doi.org/10.1093/cz/zow094).
- Ennis RS, Brandt ME, Grimes KRW, Smith TB. 2016.** Coral reef health response to chronic and acute changes in water quality in St. Thomas, United States Virgin Islands. *Marine Pollution Bulletin* **111**:418–427 DOI [10.1016/j.marpolbul.2016.07.033](https://doi.org/10.1016/j.marpolbul.2016.07.033).
- Excoffier L, Lischer HEL. 2010.** Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources* **10**:564–567 DOI [10.1111/j.1755-0998.2010.02847.x](https://doi.org/10.1111/j.1755-0998.2010.02847.x).
- Fabina NS, Putnam HM, Franklin EC, Stat M, Gates RD. 2012.** Transmission mode predicts specificity and interaction patterns in coral-*Symbiodinium* networks. *PLOS ONE* **7**:e44970 DOI [10.1371/journal.pone.0044970.s006](https://doi.org/10.1371/journal.pone.0044970.s006).
- Feder JL, Egan SP, Nosil P. 2012.** The genomics of speciation-with-gene-flow. *Trends in Genetics* **28**:342–350 DOI [10.1016/j.tig.2012.03.009](https://doi.org/10.1016/j.tig.2012.03.009).
- Fenner D. 2005.** *Corals of Hawaii*. Honolulu: Mutual Publishing, LLC.
- Flot JF. 2010.** SeqPHASE: a web tool for interconverting PHASE input/output files and FASTA sequence alignments. *Molecular Ecology Resources* **10**:162–166 DOI [10.1111/j.1755-0998.2009.02732.x](https://doi.org/10.1111/j.1755-0998.2009.02732.x).
- Forsman ZH, Barshis DJ, Hunter CL, Toonen RJ. 2009.** Shape-shifting corals: molecular markers show morphology is evolutionarily plastic in *Porites*. *BMC Evolutionary Biology* **9**:45 DOI [10.1186/1471-2148-9-45](https://doi.org/10.1186/1471-2148-9-45).
- Forsman Z, Wellington GM, Fox GE, Toonen RJ. 2015.** Clues to unraveling the coral species problem: distinguishing species from geographic variation in *Porites* across the Pacific with molecular markers and microskeletal traits. *PeerJ* **3**:e751 DOI [10.7717/peerj.751/supp-1](https://doi.org/10.7717/peerj.751/supp-1).
- Franklin EC, Jokiel PL, Donahue MJ. 2013.** Predictive modeling of coral distribution and abundance in the Hawaiian Islands. *Marine Ecology Progress Series* **481**:121–132 DOI [10.3354/meps10252](https://doi.org/10.3354/meps10252).
- Gaither MR, Szabó Z, Crepeau MW, Bird CE, Toonen RJ. 2011.** Preservation of corals in salt-saturated DMSO buffer is superior to ethanol for PCR experiments. *Coral Reefs* **30.2(2)**:329–333.
- Gorospe KD, Donahue MJ, Karl SA. 2015.** The importance of sampling design: spatial patterns and clonality in estimating the genetic diversity of coral reefs. *Marine Biology* **162.5**:917–928 DOI [10.1007/s00227-015-2634-8](https://doi.org/10.1007/s00227-015-2634-8).
- Gorospe KD, Karl SA. 2011.** Small-scale spatial analysis of in situ sea temperature throughout a single coral patch reef. *Juornal of Marine Biology* **2011**:1–12 DOI [10.1155/2011/719580](https://doi.org/10.1155/2011/719580).
- Gorospe KD, Karl SA. 2013.** Genetic relatedness does not retain spatial pattern across multiple spatial scales: dispersal and colonization in the coral, *Pocillopora damicornis*. *Molecular Ecology* **22**:3721–3736 DOI [10.1111/mec.12335](https://doi.org/10.1111/mec.12335).

- Gorospe KD, Karl SA. 2015.** Depth as an organizing force in *Pocillopora damicornis*: intra-reef genetic architecture. *PLOS ONE* **10**:e0122127–17
[DOI 10.1371/journal.pone.0122127](https://doi.org/10.1371/journal.pone.0122127).
- Graham NAJ. 2014.** Habitat complexity: coral structural loss leads to fisheries declines. *Current Biology* **24**:R359–R361 [DOI 10.1016/j.cub.2014.03.069](https://doi.org/10.1016/j.cub.2014.03.069).
- Guadayol Ò, Silbiger NJ, Donahue MJ, Thomas FIM. 2014.** Patterns in temporal variability of temperature, oxygen and pH along an environmental gradient in a coral reef. *PLOS ONE* **9**:e85213 [DOI 10.1371/journal.pone.0085213](https://doi.org/10.1371/journal.pone.0085213).
- Hogan JD, Thiessen RJ, Heath DD. 2010.** Variability in connectivity indicated by chaotic genetic patchiness within and among populations of a marine fish. *Marine Ecology Progress Series* **417**:263–275 [DOI 10.3354/meps08793](https://doi.org/10.3354/meps08793).
- Holland S. 2009.** *Analytic rarefaction, Version 2, software*. Athens: Hunt Mountain Software.
- Hughes TP, Graham NAJ, Jackson JBC, Mumby PJ, Steneck RS. 2010.** Rising to the challenge of sustaining coral reef resilience. *Trends in Ecology & Evolution* **25**:633–642 [DOI 10.1016/j.tree.2010.07.011](https://doi.org/10.1016/j.tree.2010.07.011).
- Hunter CL. 1988.** Genotypic diversity and population structure of the Hawaiian reef coral, *Porites compressa*. Ph.D. Dissertation, University of Hawaii at Manoa.
- Johnson MS, Black R. 1982.** Chaotic genetic patchiness in an intertidal limpet, *Siphonaria* sp. *Marine Biology* **70**:157–164 [DOI 10.1007/BF00397680](https://doi.org/10.1007/BF00397680).
- Johnson MS, Black R. 1984.** Pattern beneath the chaos: the effect of recruitment on genetic patchiness in an intertidal limpet. *Evolution* **38**:1371–1383
[DOI 10.1111/j.1558-5646.1984.tb05658.x](https://doi.org/10.1111/j.1558-5646.1984.tb05658.x).
- Kenkel CD, Goodbody-Gringley G, Caillaud D, Davies SW, Bartels E, Matz MV. 2013.** Evidence for a host role in thermotolerance divergence between populations of the mustard hill coral (*Porites astreoides*) from different reef environments. *Molecular Ecology* **22**:4335–4348 [DOI 10.1111/mec.12391](https://doi.org/10.1111/mec.12391).
- LaJeunesse T, Thornhill D, Cox E, Stanton F, Fitt W, Schmidt G. 2004.** High diversity and host specificity observed among symbiotic dinoflagellates in reef coral communities from Hawaii. *Coral Reefs* **23**:596–603 [DOI 10.1007/s00338-004-0428-4](https://doi.org/10.1007/s00338-004-0428-4).
- Levas SJ, Grottoli AG, Hughes A, Osburn CL, Matsui Y. 2013.** Physiological and Biogeochemical traits of bleaching and recovery in the mounding species of coral *Porites lobata*: implications for resilience in mounding corals. *PLOS ONE* **8**:e63267
[DOI 10.1371/journal.pone.0063267.t006](https://doi.org/10.1371/journal.pone.0063267.t006).
- McRae BH, Beier P. 2007.** Circuit theory predicts gene flow in plant and animal populations. *Proceedings of the National Academy of Sciences of the United States of America* **104**:19885–19890 [DOI 10.1073/pnas.0706568104](https://doi.org/10.1073/pnas.0706568104).
- Morgan KM, Perry CT, Smithers SG, Johnson JA, Daniell JJ. 2016.** Evidence of extensive reef development and high coral cover in nearshore environments: implications for understanding coral adaptation in turbid settings. *Scientific Reports* **6**:29616
[DOI 10.1038/srep29616](https://doi.org/10.1038/srep29616).

- Nosil P, Funk DJ, Ortiz-Barrientos D. 2009. Divergent selection and heterogeneous genomic divergence. *Molecular Ecology* **18**:375–402
DOI [10.1111/j.1365-294X.2008.03946.x](https://doi.org/10.1111/j.1365-294X.2008.03946.x).
- Orsini L, Vanoverbeke J, Swillen I, Mergeay J, De Meester L. 2013. Drivers of population genetic differentiation in the wild: isolation by dispersal limitation, isolation by adaptation and isolation by colonization. *Molecular Ecology* **22**:5983–5999
DOI [10.1111/mec.12561](https://doi.org/10.1111/mec.12561).
- Polato NR, Concepcion GT, Toonen RJ, Baums IB. 2010. Isolation by distance across the Hawaiian Archipelago in the reef-building coral *Porites lobata*. *Molecular Ecology* **19**:4661–4677 DOI [10.1111/j.1365-294X.2010.04836](https://doi.org/10.1111/j.1365-294X.2010.04836).
- Presto KM, Storlazzi CD, Logan JB, Reiss TE, Rosenberger KJ. 2012. Coastal circulation and potential coral-larval dispersal in Maunaloa Bay, Oahu, Hawaii—measurements of waves, currents, temperature, and salinity June-2010. U.S. Geological Survey Open-File Report 2012-1040.
- Puritz JB, Toonen RJ. 2011. Coastal pollution limits pelagic larval dispersal. *Nature Communications* **2**:1–8 DOI [10.1038/ncomms1238](https://doi.org/10.1038/ncomms1238).
- R Core Team. 2014. *R: a language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing.
- Richmond RH. 2009. HCRI Project Progress Report FY2008: Watersheds impacts on coral reefs in Maunaloa Bay, Oahu, Hawaii.
- Richmond RH, Wolanski E. 2011. Coral research: past efforts and future horizons. In: Dubinsky Z, Stambler N, eds. *Coral reefs an ecosystem in transition*. Dordrecht: Springer, 3–12.
- Rodgers KS, Jokiel PL, Brown EK, Hau S, Sparks R. 2015. Over a decade of change in spatial and temporal dynamics of Hawaiian coral reef communities 1. *Pacific Science* **69**:1–13 DOI [10.2984/69.1.1](https://doi.org/10.2984/69.1.1).
- Roff G, Bejarano S, Bozec Y-M, Nugues M, Steneck RS, Mumby PJ. 2014. *Porites* and the Phoenix effect: unprecedented recovery after a mass coral bleaching event at Rangiroa Atoll, French Polynesia. *Marine Biology* **161**:1385–1393
DOI [10.1007/s00227-014-2426-6](https://doi.org/10.1007/s00227-014-2426-6).
- Rundle HD, Nosil P. 2005. Ecological speciation. *Evolution* **8**:336–352
DOI [10.1111/j.1461-0248.2004.00715.x](https://doi.org/10.1111/j.1461-0248.2004.00715.x).
- Schweinsberg M, Tollrian R, Lampert KP. 2016. Genetic variation in the massive coral *Porites lobata*. *Marine Biology* **163**:1–12 DOI [10.1007/s00227-016-3022-8](https://doi.org/10.1007/s00227-016-3022-8).
- Selkoe KA, D'Aloia CC, Crandall ED, Iacchi M, Liggins L, Puritz JB, Heyden von der S, Toonen RJ. 2016. A decade of seascape genetics: contributions to basic and applied marine connectivity. *Marine Ecology Progress Series* **554**:1–19
DOI [10.3354/meps11792](https://doi.org/10.3354/meps11792).
- Shearer TL, Van Oppen MJH, Romano SL, Wörheide G. 2002. Slow mitochondrial DNA sequence evolution in the Anthozoa (Cnidaria). *Molecular Ecology* **11**:2475–2487
DOI [10.1046/j.1365-294X.2002.01652.x](https://doi.org/10.1046/j.1365-294X.2002.01652.x).
- Slatkin M. 1993. Isolation by distance in equilibrium and non-equilibrium populations. *Evolution* **47**(1):264–279 DOI [10.1111/j.1558-5646.1993.tb01215.x](https://doi.org/10.1111/j.1558-5646.1993.tb01215.x).

- Smith TB, Nemeth RS, Blondeau J, Calnan JM, Kadison E, Herzlieb S. 2008b.** Assessing coral reef health across onshore to offshore stress gradients in the US Virgin Islands. *Marine Pollution Bulletin* **56**:1983–1991 DOI [10.1016/j.marpolbul.2008.08.015](https://doi.org/10.1016/j.marpolbul.2008.08.015).
- Smith LW, Wirshing HH, Baker AC, Birkeland C. 2008a.** Environmental versus genetic influences on growth rates of the corals *Pocillopora eydouxi* and *Porites lobata* (Anthozoa: Scleractinia) 1. *Pacific Science* **62**:57–69 DOI [10.2984/1534-6188\(2008\)62\[57:EVGIOG\]2.0.CO;2](https://doi.org/10.2984/1534-6188(2008)62[57:EVGIOG]2.0.CO;2).
- Stafford-Smith MG. 1993.** Sediment-rejection efficiency of 22 species of Australian scleractinian corals. *Marine Biology* **115**:229–243 DOI [10.1007/BF00346340](https://doi.org/10.1007/BF00346340).
- Stephens M, Smith NJ, Donnelly P. 2001.** A new statistical method for haplotype reconstruction from population data. *American Journal of Human Genetics* **68**:978–989 DOI [10.1086/319501](https://doi.org/10.1086/319501).
- Storlazzi CD, Field ME. 2008.** Winds, waves, tides, and the resulting flow patterns and fluxes of water, sediment, and coral larvae off West Maui, Hawaii. USGS Open-File Report 2008-1215.
- Storlazzi CD, McManus MA, Logan JB, McLaughlin BE. 2006.** Cross-shore velocity shear, eddies and heterogeneity in water column properties over fringing coral reefs: West Maui, Hawaii. *Continental Shelf Research* **26**:401–421 DOI [10.1016/j.csr.2005.12.006](https://doi.org/10.1016/j.csr.2005.12.006).
- Storlazzi CD, Presto KM, Logan JB, Field ME. 2010.** Coastal circulation and sediment dynamics in Maunaloa Bay, Oahu, Hawaii. USGS Open-File Report 2010-1217.
- Thomas L, Kennington WJ, Stat M, Wilkinson SP, Kool JT, Kendrick GA. 2015.** Isolation by resistance across a complex coral reef seascape. *Proceedings of the Royal Society B: Biological Sciences* **282**:20151217 DOI [10.1098/rspb.2015.1217](https://doi.org/10.1098/rspb.2015.1217).
- Thompson A, Schroeder T, Brando VE, Schaffelke B. 2014.** Coral community responses to declining water quality: Whitsunday Islands, Great Barrier Reef, Australia. *Coral Reefs* **33**:923–928 DOI [10.1007/s00338-014-1201-y](https://doi.org/10.1007/s00338-014-1201-y).
- Tisthammer KH, Forsman ZH, Sindorf VL, Massey TL, Bielecki CR, Toonen RJ. 2016.** The complete mitochondrial genome of the lobe coral *Porites lobata* (Anthozoa: Scleractinia) sequenced using ezRAD. *Mitochondrial DNA Part B* **1**:247–249 DOI [10.1080/23802359.2016.1157770](https://doi.org/10.1080/23802359.2016.1157770).
- Tisthammer KH, Richmond RH. 2018.** Corallite skeletal morphological variation in Hawaiian *Porites lobata*. *Coral Reefs* **65**:560–514 DOI [10.1007/s00338-018-1670-5](https://doi.org/10.1007/s00338-018-1670-5).
- Toonen RJ, Grosberg RK. 2011.** Causes of chaos: spatial and temporal genetic heterogeneity in the intertidal anomuran crab *Petrolisthes cinctipes*. *Phylogeography and Population Genetics in Crustacea* 75–107.
- Van Dam JW, Negri AP, Uthicke S, Mueller JF. 2011.** Chemical pollution on coral reefs: exposure and ecological effects. In: Sánchez-Bayo F, van den Brink PJ, Mann RM, eds. *Ecological impact of toxic chemicals*. Bussum, The Netherlands: Bentham Science Publishers, 187–211.
- Vargas-Angel B. 2017.** Maui_Baseline_report_2017. 1–49 DOI [10.7289/V5/SP-PIFSC-17-001](https://doi.org/10.7289/V5/SP-PIFSC-17-001).

- Veron JEN. 1995.** *Corals in space and time: the biogeography and evolution of the Scleractinia*. Ithaca: Cornell University Press.
- Veron JEN. 2000.** *Corals of the world*. Townsville: Australian Institute of Marine Science.
- Veron JEN, Pichon M. 1982.** *Scleractinia of Eastern Australia part IV family Poritidae*. Canberra: Australian National University Press.
- Vollmer SV, Palumbi SR. 2002.** Hybridization and the evolution of reef coral diversity. *Science* **296**:2023–2025 DOI [10.1126/science.1069524](https://doi.org/10.1126/science.1069524).
- Wang IJ, Bradburd GS. 2014.** Isolation by environment. *Molecular Ecology* **23**:5649–5662 DOI [10.1111/mec.12938](https://doi.org/10.1111/mec.12938).
- Wenger AS, Williamson DH, Da Silva ET, Ceccarelli DM, Browne NK, Petus C, Devlin MJ. 2015.** Effects of reduced water quality on coral reefs in and out of no-take marine reserves. *Conservation Biology* **30**:142–153 DOI [10.1111/cobi.12576](https://doi.org/10.1111/cobi.12576).
- Williams I, Vargas B, White D, Callender T. 2014.** West Maui Wahikuli & Honokowai Priority Watershed Area: Reef Condition Report. Scientific Report by Hawaii Division of Aquatic Resources, Ridge 2 Reef Initiative, NOAA Coral Reef Conservation Program.
- Wolanski E, Martinez JA, Richmond RH. 2009.** Quantifying the impact of watershed urbanization on a coral reef: Maunalua Bay, Hawaii. *Estuarine, Coastal and Shelf Science* **84**:259–268 DOI [10.1016/j.ecss.2009.06.029](https://doi.org/10.1016/j.ecss.2009.06.029).
- Wright S. 1943.** Isolation by distance. *Genetics* **28**:114–138.